

Microscopic dynamics of an amorphous C₆₀x/C₇₀(1 – x) fullerene mixture

Khusnutdinoff R., Mokshin A., Takhaviev I.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2015, Pleiades Publishing, Ltd. This paper presents the results of investigation of the propagation mechanism of collective excitations in amorphous C₆₀x/C₇₀(1 – x) fullerene mixtures (with equimolar concentration $x = 0.50$), which were obtained using molecular dynamics simulation. The critical glass-transition temperature of the system $T_c = 1548$ K was determined from the change in the behavior of the Wendt-Abraham parameter. Spectral densities of the time correlation functions of the longitudinal (Formula presented.) and transverse (Formula presented.) currents for a wide region of wave numbers at temperatures below the glass-transition temperature were calculated. It was found that the dynamics of density fluctuations in amorphous C₆₀x/C₇₀(1 – x) fullerene mixtures is characterized by two dispersion acoustic-like branches of the longitudinal and transverse polarizations. The influence of poly-dispersity and form factor of the molecule of the mixture component on the microscopic dynamics of the density fluctuation in multicomponent systems was established.

<http://dx.doi.org/10.1134/S1063783415020183>
